Composing Cardinal Direction Relations

Spiros Skiadopoulos[1] and Manolis Koubarakis[2]

[1] Dept. of Electrical and Computer Engineering
National Technical University of Athens
Zographou 157 73 Athens, Greece
spiros@dblab.ntua.gr, www.dblab.ntua.gr/~spiros

[2] Dept. of Electronic and Computer Engineering
Technical University of Crete
Chania 73100 Crete, Greece
manolis@ced.tuc.gr, www.ced.tuc.gr/~manolis

Abstract. We study the recent proposal of Goyal and Egenhofer who presented a model for qualitative spatial reasoning about cardinal directions. Our approach is formal and complements the presentation of Goyal and Egenhofer. We focus our efforts on the operation of composition for two cardinal direction relations. We point out that the only published method to compute the composition does not always work correctly. Then we consider progressively more expressive classes of cardinal direction relations and give composition algorithms for these classes. Our theoretical framework allows us to prove formally that our algorithms are correct. Finally, we demonstrate that in some cases, the binary relation resulting from the composition of two cardinal direction relations cannot be expressed using the relations defined by Goyal and Egenhofer.

1 Introduction

Qualitative spatial reasoning has received a lot of attention in the areas of Geographic Information Systems [3, 5], Artificial Intelligence [2, 4, 16, 17], Databases [14] and Multimedia [18]. Several kinds of useful spatial relations have been studied so far e.g., topological relations [1–3, 16, 17], cardinal direction relations [7, 10] and qualitative distance relations [5, 20].

In this paper we concentrate on *cardinal direction relations* [7, 10, 12]. Cardinal direction relations describe how regions of space are placed relative to one another (e.g., region a is *north of* region b). Our starting point is the recent model of Goyal and Egenhofer [6, 7]. This model is currently one of the most cognitively plausible models for qualitative reasoning with cardinal directions. It works with extended regions and has potential in Multimedia and Geographic Information Systems applications.

In a recent paper [19] we have studied the problem of checking the consistency of a given set of constraints in the model of [6, 7] and its variations. In this paper we concentrate on the problem of *composing* cardinal direction relations.

The composition operation for various kinds of spatial relations has received a lot of attention in the literature [3, 5, 12]. *Composition tables* are typically used as a mechanism for inferring new spatial relations from existing ones. This is very helpful in the following situations:

1. Detecting possible inconsistencies in a given set of spatial relations [10, 17].
2. Preprocessing spatial queries so that inconsistent queries are detected or the search space is pruned [13].

Therefore it is very important to study the composition operation and define appropriate composition tables for interesting models of spatial relations such as [6, 7].

The technical contributions of this paper can be summarized as follows:

- We give formal definitions for the cardinal direction relations that can be expressed in the model of Goyal and Egenhofer [7].
- We use our formal framework to study the composition operation for cardinal direction relations in the model of [7]. Goyal and Egenhofer first studied this operation in [7] but their method does not always work correctly.
- The previous observation leaves us with the task of finding a correct method for computing the composition. To do this, we consider progressively more expressive classes of cardinal direction relations and give composition algorithms for these classes. Our theoretical framework allows us to *prove formally* that our algorithms are correct.
  Finally, when we consider the problem in its generality, we show something very interesting: the binary relation resulting from the composition of some cardinal direction relations *cannot even be expressed* using the relations defined in [7] unless the language is augmented by extra predicates (Section 5).

The rest of the paper is organized as follows. Section 2 presents the formal model. In Sections 3 and 4 we consider two subclasses of cardinal direction relations and give composition algorithms for these classes. In Section 5 we show that the result of the composition of some cardinal direction relations cannot be expressed using the relations defined in [7]. Our conclusions are presented in Section 6.

2 A Formal Model for Cardinal Direction Information

We consider the Euclidean space $\Re^2$. *Regions* are defined as non-empty sets of points in $\Re^2$. A region a is *connected* if for every pair of points in a there exists a line (not necessarily a straight one) joining the two points such that all the points on the line are also in a. The connectedness property prevents a region from having disjoint parts. A region is *disconnected* if it is not connected.

Let a be a region. The *greatest lower bound* or the *infimum* [11] of the *projection* of region a on the x-axis (respectively y-axis) is denoted by $inf_x(a)$ (respectively $inf_y(a)$). The *least upper bound* or the *supremum* of the *projection*



Fig. 1. A region and its bounding box

of region a on the x-axis (respectively y-axis) is denoted by $sup_x(a)$ (respectively $sup_y(a)$).

The *minimum bounding box* of a region a, denoted by $mbb(a)$, is the rectangular region formed by the straight lines $x = inf_x(a)$, $x = sup_x(a)$, $y = inf_y(a)$ and $y = sup_y(a)$ (see Figure 1). Obviously, the projections on the x-axis (respectively y-axis) of a region and its minimum bounding box have the same endpoints.

We will consider throughout the paper the following types of regions:

- Regions that are connected, non-empty subsets of $\Re^2$ and homeomorphic images of two-dimensional disks [7, 15]. The set of these regions will be denoted by REG^{disk}.
- Regions that are formed by finite unions of regions in REG^{disk}. The set of these regions will be denoted by REG^{disk*}. Notice that regions in REG^{disk*} can be disconnected.



Fig. 2. Regions

In Figure 2, regions a, b_1, b_2 and b_3 are in REG^{disk} (also in REG^{disk*}) and region $b = b_1 \cup b_2 \cup b_3$ is in REG^{disk*}.

Regions in REG^{disk} have been previously studied in [7, 15]. They can be used to model areas in various interesting applications, e.g., land parcels in Geographic Information Systems [3, 5]. In the sequel we will formally define cardinal direction relations for regions in REG^{disk}. To this end, we will need regions in REG^{disk*}.

Let us now consider two arbitrary regions a and b in REG^{disk}. Let region a be related to region b through a cardinal direction relation (e.g., a is north of b). Region b will be called the *reference* region (i.e., the region *to* which the relation is described) while region a will be called the *primary* region (i.e., the region *from* which the relation is described) [7]. The axes forming the minimum bounding box of the reference region b divide the space into 9 tiles (Figure 3a). The peripheral tiles correspond to the eight cardinal direction relations *south*, *southwest*, *west*, *northwest*, *north*, *northeast*, *east* and *southeast*. These tiles will be denoted by $S(b)$, $SW(b)$, $W(b)$, $NW(b)$, $N(b)$, $NE(b)$, $E(b)$ and $SE(b)$ respectively. The central area corresponds to the region's minimum bounding box and is denoted by $B(b)$. By definition each one of these tiles includes the parts of the axes forming it. The union of all 9 tiles is $\Re^2$.



Fig. 3. Reference tiles and relations

If a primary region a is included (in the set-theoretic sense) in tile $S(b)$ of some reference region b (Figure 3b) then we say that *a is south of b* and we write $a\ S\ b$. Similarly, we can define *southwest* (SW), *west* (W), *northwest* (NW), *north* (N), *northeast* (NE), *east* (E), *southeast* (SE) and *bounding box* (B) relations.

If a primary region a lies partly in the area $NE(b)$ and partly in the area $E(b)$ of some reference region b (Figure 3c) then we say that *a is partly northeast and partly east of b* and we write $a\ NE{:}E\ b$.

The general definition of a cardinal direction relation is as follows.

Definition 1. *An* atomic cardinal direction relation *is an element of the set* $\{B, S, SW, W, NW, N, NE, E, SE\}$. *A* basic cardinal direction relation *is an atomic cardinal direction relation or an expression* $R_1{:}\cdots{:}R_k$ *where* $2 \leq k \leq 9$, $R_1, \ldots, R_k \in \{B, S, SW, W, NW, N, NE, E, SE\}$, $R_i \neq R_j$ *for every* i, j *such that* $1 \leq i, j \leq k$ *and* $i \neq j$, *and the tiles* $R_1(b), \ldots, R_k(b)$ *form a region of* REG^{disk} *for any reference region b.*

Example 1. The following are basic cardinal direction relations:

$$S,\quad NE{:}E \quad \text{and} \quad B{:}S{:}SW{:}W{:}NW{:}N{:}E{:}SE.$$

Regions involved in these relations are shown in Figures 3b, 3c and 3d respectively.